

## Titchmarsh's Theorem for the Bessel Transform

**R. Daher, M. El Hamma and A. El Houasni**

Department of Mathematics, Faculty of Sciences Ain Chock  
 University of Hassan II, Casablanca, Morocco  
 e-mail: elmohamed77@yahoo.fr

**Abstract** In [1] Titchmarsh proved some theorems on the classical Fourier transform of functions satisfying conditions related to the Cauchy-Lipschitz conditions on the Euclidean space  $\mathbb{R}$ . In this paper we extend one those theorems for the Bessel transform for function on half-line  $[0, \infty)$  in a weighted  $L_p$ -metric are studied with the use of Bessel generalized translation.

**Keywords** Bessel operator, Bessel transform, Bessel generalized translation.

**2010 Mathematics Subject Classification** 42B10, 42A38, 42B37.

### 1 Introduction and Preliminaries

Integral transforms and their inverses (e.g., the Fourier-Bessel transform) are widely used to solve various problems in calculus, mechanics, mathematical physics, and computational mathematics (see, e.g., [1, 7]).

In this paper, we use the Bessel generalized translation, it is one of the most important generalized translations on the half-line  $\mathbb{R}^+ = [0, +\infty)$  [4, 5]. The Bessel generalized translation is used while studying various problems connected with Bessel operators (see [3, 8]).

Titchmarsh ([1], Theorem 84) characterized the set of functions in  $L^p(\mathbb{R})$  satisfying the estimate, namely we have

**Theorem 1.1** *Let  $f(x)$  belong to  $L^p(\mathbb{R})$  ( $1 < p \leq 2$ ), and let*

$$\int_{-\infty}^{+\infty} |f(x+h) - f(x-h)|^p dx = O(h^{\alpha p}), \quad (0 < \alpha \leq 1)$$

*as  $h \rightarrow 0$ . Then  $\mathcal{F}(f)(x)$  belongs to  $L^\beta(\mathbb{R})$  for*

$$\frac{p}{p + \alpha p - 1} < \beta \leq \frac{p}{p - 1}$$

*where  $\mathcal{F}(f)$  stands for the Fourier transform of  $f$*

The main of this paper is to establish an analog of Theorem 1.1 in the Bessel operators setting by means of the Bessel generalized translation.

Let

$$B = \frac{d^2}{dt^2} + \frac{(2\alpha + 1)}{t} \frac{d}{dt},$$

be the Bessel differential operator. By  $j_\alpha(t)$  denote the Bessel normed function of the first kind, i.e

$$j_\alpha(t) = \frac{2^\alpha \Gamma(\alpha + 1) J_\alpha(t)}{t^\alpha},$$

where  $J_\alpha(x)$  is the Bessel function of the first kind and  $\Gamma(x)$  is the gamma-function (see[4]). The function  $y = j_\alpha(t)$  satisfies the differential equation  $By + y = 0$  with the initial conditions  $y(0) = 1$  and  $y'(0) = 0$ . The function  $j_\alpha(t)$  is infinitely differentiable, entire analytic.

Assume that  $L_\alpha^p(\mathbb{R}_+)$ ,  $\alpha > -\frac{1}{2}$  and  $1 < p \leq 2$ , is the Banach space of mesurable functions  $f(t)$  on  $\mathbb{R}_+$  with the finite norm

$$\|f\| = \|f\|_{p,\alpha} = \left( \int_0^\infty |f(t)|^p t^{2\alpha+1} dt \right)^{1/p}.$$

In  $L_\alpha^p(\mathbb{R}_+)$ , consider the Bessel generalized translation  $T_h$  (see [3, p.121])

$$T_h f(x) = \frac{\Gamma(\alpha+1)}{\Gamma(1/2)\Gamma(\alpha+\frac{1}{2})} \int_0^\pi f(\sqrt{x^2+h^2-2xh\cos t}) \sin^{2\alpha} t dt, \quad \alpha > -\frac{1}{2}, \quad 0 \leq h \leq 1$$

which corresponds to the Bessel operator  $B$ .

It is easy to see that

$$T_0 f(x) = f(x)$$

If  $f(x)$  has a continuous first derivative, then

$$\frac{\partial}{\partial h} T_h f(x)|_{h=0} = 0.$$

If it has a continuous second derivative, then  $u(x, h) = T_h f(x)$  solves the Cauchy problem

$$\frac{\partial^2 u}{\partial x^2} + \frac{2\alpha+1}{x} \frac{\partial u}{\partial x} = \frac{\partial^2 u}{\partial h^2} + \frac{2\alpha+1}{h} \frac{\partial u}{\partial h}$$

and

$$u|_{h=0} = f(x), \quad \frac{\partial u}{\partial h}|_{h=0} = 0.$$

The operator  $T_h$  is linear, homogeneous, and continuous. Below are some properties of this operator (see [3, pp. 124-125]):

$$(i) \quad T_h j_\alpha(\lambda x) = j_\alpha(\lambda h) j_\alpha(\lambda x)$$

(ii)  $T_h$  is self-adjoint. If  $f(x)$  is continuous function such that  $\int_0^\infty x^{2\alpha+1} |f(x)| dx < \infty$  and  $g(x)$  is continuous and bounded for all  $x \geq 0$ , then

$$\int_0^\infty (T_h f(x)) g(x) x^{2\alpha+1} dx = \int_0^\infty f(x) (T_h g(x)) x^{2\alpha+1} dx.$$

$$(iii) \quad T_h f(x) = T_x f(h).$$

$$(iv) \quad \|T_h f - f\| \longrightarrow 0 \text{ as } h \longrightarrow 0.$$

The Bessel transform defined by the formula (see [3, 4, 6])

$$\widehat{f}(\lambda) = \int_0^\infty f(t) j_\alpha(\lambda t) t^{2\alpha+1} dt; \quad \lambda \in \mathbb{R}_+$$

The inverse Bessel transform is given by the formula

$$f(t) = (2^\alpha \Gamma(\alpha + 1))^{-2} \int_0^\infty \widehat{f}(\lambda) j_\alpha(\lambda t) \lambda^{2\alpha+1} d\lambda$$

The following relation connect the Bessel generalized translation, and the Bessel transform in [2], we have

$$(\widehat{T_h f})(\lambda) = j_\alpha(\lambda h) \widehat{f}(\lambda) \quad (1)$$

For  $\alpha > -\frac{1}{2}$ , we introduce the Bessel normalized function of the first kind  $j_\alpha$  defined by

$$j_\alpha(x) = \Gamma(\alpha + 1) \sum_{n=0}^\infty \frac{(-1)^n (x/2)^{2n}}{n! \Gamma(n + \alpha + 1)} \quad (2)$$

Moreover, from (2) we see that

$$\lim_{x \rightarrow 0} \frac{(j_\alpha(x) - 1)}{x^2} \neq 0$$

by consequence, there exist  $C > 0$  and  $\eta > 0$  satisfying

$$|x| \leq \eta \implies |j_\alpha(x) - 1| \geq C|x|^2 \quad (3)$$

## 2 An Analog of Titchmarsh's Theorem

In this section, we give an analog of Titchmarsh's Theorem 84 for the Bessel transform.

**Theorem 2.1** *Let  $f(x)$  belong to  $L_\alpha^p(\mathbb{R}_+)$ , ( $1 < p \leq 2$ ), and let*

$$\int_0^\infty |T_h f(x) - f(x)|^p x^{2\alpha+1} dx = O(h^{\gamma p}), \quad (0 < \gamma \leq 2)$$

*as  $h \rightarrow 0$ . Then  $\widehat{f}(x)$  belongs to  $L_\alpha^\beta(\mathbb{R}_+)$ , for*

$$\frac{2p\alpha + 2p}{2p + 2\alpha(p - 1) + \gamma p - 2} < \beta \leq \frac{p}{p - 1}$$

**Proof:** For a fixed  $h$  the Bessel transform of  $T_h f(x)$  is  $j_\alpha(hx)\widehat{f}(x)$ . Hence the Bessel transform of  $T_h f(x) - f(x)$ , as a function of  $x$ , is  $(j_\alpha(hx) - 1)\widehat{f}(x)$ .

Hence

$$\begin{aligned} \int_0^\infty |j_\alpha(hx) - 1|^{p'} |\widehat{f}(x)|^{p'} x^{2\alpha+1} dx &< K(p) \left( \int_0^\infty |T_h f(x) - f(x)|^p x^{2\alpha+1} dx \right)^{1/p-1} \\ &< K(p) h^{\gamma p'} \end{aligned}$$

From formula (3), we have

$$\int_0^{\eta/h} |hx|^{2p'} |\widehat{f}(x)|^{p'} x^{2\alpha+1} dx < K(p) h^{\gamma p'}$$

Then

$$\int_0^{\eta/h} x^{2p'} |\widehat{f}(x)|^{p'} x^{2\alpha+1} dx < K(p) h^{(\gamma-2)p'}$$

let

$$\phi(\xi) = \int_1^\xi |x^2 \widehat{f}(x)|^{\beta} x^{(2\alpha+1)\frac{p'}{\beta}} dx$$

Then, if  $\beta < p'$

$$\begin{aligned} \phi(\xi) &\leq \left( \int_1^\xi |x^2 \widehat{f}(x)|^{p'} x^{2\alpha+1} dx \right)^{\beta/p'} \left( \int_1^\xi dx \right)^{1-\beta/p'} \\ &= O(\xi^{(2-\gamma)p'\frac{\beta}{p'}\xi^{1-\frac{\beta}{p'}}}) \\ &= O(\xi^{2\beta-\gamma\beta+1-\frac{\beta}{p'}}) \end{aligned}$$

Hence

$$\begin{aligned} \int_1^\xi |\widehat{f}(\xi)|^\beta x^{2\alpha+1} dx &= \int_1^\xi x^{-2\beta-(2\alpha+1)\frac{\beta}{p'}} \phi'(x) x^{2\alpha+1} dx \\ &= \xi^{-2\beta-(2\alpha+1)\frac{\beta}{p'}} \xi^{2\alpha+1} \phi(\xi) + (2\beta + (2\alpha+1)\frac{\beta}{p'} - (2\alpha+1)) \int_1^\xi x^{-2\beta-(2\alpha+1)\frac{\beta}{p'}+2\alpha} \phi(x) dx \\ &= O(\xi^{-2\beta-(2\alpha+1)\frac{\beta}{p'}+2\alpha+1+1-\gamma\beta+\beta(\frac{p+1}{p})}) + O\left(\int_1^\infty x^{-2\beta-(2\alpha+1)\frac{\beta}{p'}+2\alpha} x^{1-\gamma\beta+\beta(\frac{p+1}{p})} dx\right) \\ &= O(\xi^{-2\beta-(2\alpha+1)\frac{\beta}{p'}+2\alpha+2-\gamma\beta+\beta(\frac{p+1}{p})}) \end{aligned}$$

and this bounded as  $\xi \rightarrow \infty$  if  $-2\beta - (2\alpha+1)\frac{\beta}{p'} + 2\alpha + 2 - \gamma\beta + \beta(\frac{p+1}{p}) < 0$  i.e, if

$$\beta > \frac{2p\alpha + 2p}{2p + 2\alpha(p-1) + \gamma p - 2}$$

This proves the theorem.

### 3 Conclusion

There are many theorems known about to classical Fourier transform can be generalized for the Bessel transform, among them Titchmarsh's Theorem. In this work we have succeeded to generalise this theorem for the Bessel transform in the space  $L_\alpha^p(\mathbb{R}_+)$ . We proved that if  $f(x)$  belongs to  $L_\alpha^p(\mathbb{R}_+)$ , and that

$$\int_0^\infty |T_h f(x) - f(x)|^p x^{2\alpha+1} dx = O(h^\gamma), \quad (0 < \gamma \leq 2)$$

as  $h \rightarrow 0$ , then its Bessel transform  $\hat{f}$  belongs to  $L_\alpha^\beta(\mathbb{R}_+)$  for

$$\frac{2p\alpha + 2p}{2p + 2\alpha(p-1) + \gamma p - 2} < \beta \leq \frac{p}{p-1},$$

where  $0 < \gamma < 1$  and  $1 < p \leq 2$ .

### References

- [1] Titchmarsh, E. C. (2005) *Introduction to the Theory of Fourier Integrals*. Moscow: Clarendon. Oxford. 1948, Komkniga, Moscow.
- [2] Abilov, V. A., Abilova, F. V. and Kerimov, M. K. On Estimates for the Fourier-Bessel Integral Transform in the  $L_2(\mathbb{R}_+)$ . *Zh. Vychis. Mat. Mat. Fiz.* 2009. 49(7): 1158-1166.
- [3] Kipriyanov, I.A. *Singular Elliptic Boundary Value Problems*. Moscow: Nauka. 1997.
- [4] Levitan, B.M. Expansion in Fourier Series and Integrals over Bessel Functions. *Uspekhi. Mat. Nauk.* 1951. 6(2): 102-148.
- [5] Trimèche, K. *Generalized Harmonic Analysis and Wavelet Packets*. Amsterdam: Gordon and Breach Science Publ. 2000.
- [6] Trimèche, K. Transmutation operators and mean-periodic functions associated with differential operators. *Math. Rep.* 1988. 4(1): 1-282.
- [7] Zayed, A. L. *Handbook of Function and Generalized Function Transformations*. CRC. Boca Raton. 1996.
- [8] Zhitomirskii, Y. A. Cauchy's problem for systems of linear partial differential equations with differential operators of Bessel type. *Mat. Sb.* 1955. 36(2): 299-310.